



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2010

A new crystal modification of diammonium hydrogen phosphate, (NH₄)₂(HPO₄)

Kunz, P C ; Wetzler, C ; Spingler, B

Abstract: The addition of hexafluoridophosphate salts (ammonium, silver, thallium or potassium) is usually used to precipitate complex cations from aqueous solutions. It has long been known that PF₆⁻ is sensitive towards hydrolysis under acidic conditions Gebala Jones (1969). J. Inorg. Nucl. Chem. 31, 771-776; Plakhotnyk et al. (2005). J. Fluorine Chem. 126, 27-31]. During the course of our investigation into coinage metal complexes of diphosphine ligands, we used ammonium hexafluoridophosphate in order to crystallize Ag(diphosphine)₂]PF₆ complexes. From these solutions we always obtained needle-like crystals which turned out to be the title compound, 2NH₄⁺center dot HPO₄²⁻. It was received as the hydrolysis product of NH₄PF₆. The crystals are a new modification of diammonium hydrogen phosphate. In contrast to the previously published polymorph Khan et al. (1972). Acta Cryst. B28, 2065 -2069], Z' of the title compound is 2. In the new modification of the title compound, there are eight molecules of (NH₄)₂(HPO₄) in the unit cell. The structure consists of PO₃OH and NH₄ tetrahedra, held together by O-H center dot center dot center dot O and N-H center dot center dot center dot O hydrogen bonds.

DOI: <https://doi.org/10.1107/S1600536810009839>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-44992>

Journal Article

Originally published at:

Kunz, P C; Wetzler, C; Spingler, B (2010). A new crystal modification of diammonium hydrogen phosphate, (NH₄)₂(HPO₄). Acta Crystallographica. Section E, Structure Reports Online, E66(Pt. 4):i26-i27.


DOI: <https://doi.org/10.1107/S1600536810009839>

A new crystal modification of diammonium hydrogen phosphate, $(\text{NH}_4)_2(\text{HPO}_4)$

Peter C. Kunz, Corinna Wetzel and Bernhard Spingler

Acta Cryst. (2010). **E66**, i26–i27

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



Acta Crystallographica Section E: Structure Reports Online is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. In 2007, the journal published over 5000 structures. The average publication time is less than one month.

Crystallography Journals **Online** is available from journals.iucr.org

A new crystal modification of diammonium hydrogen phosphate, (NH₄)₂(HPO₄)

Peter C. Kunz,^{a*} Corinna Wetzel^a and Bernhard Spingler^b

^aHeinrich-Heine-Universität Düsseldorf, Institut für Anorganische Chemie und Strukturchemie I, Universitätsstrasse 1, D-40225 Düsseldorf, Germany, and

^bAnorganisch-Chemisches Institut, Universität Zürich-Irchel, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland

Correspondence e-mail: peter.kunz@uni-duesseldorf.de

Received 8 February 2010; accepted 16 March 2010

Key indicators: single-crystal X-ray study; $T = 183$ K; mean $\sigma(\text{P}-\text{O}) = 0.002$ Å; R factor = 0.048; wR factor = 0.158; data-to-parameter ratio = 29.7.

The addition of hexafluoridophosphate salts (ammonium, silver, thallium or potassium) is usually used to precipitate complex cations from aqueous solutions. It has long been known that PF_6^- is sensitive towards hydrolysis under acidic conditions [Gebala & Jones (1969). *J. Inorg. Nucl. Chem.* **31**, 771–776; Plakhotnyk *et al.* (2005). *J. Fluorine Chem.* **126**, 27–31]. During the course of our investigation into coinage metal complexes of diphosphine ligands, we used ammonium hexafluoridophosphate in order to crystallize $[\text{Ag}(\text{diphosphine})_2]\text{PF}_6$ complexes. From these solutions we always obtained needle-like crystals which turned out to be the title compound, $2\text{NH}_4^+\cdot\text{HPO}_4^{2-}$. It was received as the hydrolysis product of NH_4PF_6 . The crystals are a new modification of diammonium hydrogen phosphate. In contrast to the previously published polymorph [Khan *et al.* (1972). *Acta Cryst. B* **28**, 2065–2069], Z' of the title compound is 2. In the new modification of the title compound, there are eight molecules of $(\text{NH}_4)_2(\text{HPO}_4)$ in the unit cell. The structure consists of PO_3OH and NH_4 tetrahedra, held together by $\text{O}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For the study of another crystal modification of the title compound, see: Khan *et al.* (1972). For the hydrolysis of hexafluoridophosphates, see: Akbayeva *et al.* (2006); Deifel *et al.* (2008); Fernandez-Galan *et al.* (1994); Gebala & Jones (1969); Nikitenko *et al.* (2007); Plakhotnyk *et al.* (2005).

Experimental

Crystal data

$2\text{NH}_4^+\cdot\text{HPO}_4^{2-}$

$M_r = 132.06$

Monoclinic, $P2_1/c$

$a = 11.2868$ (3) Å

$b = 15.3466$ (4) Å

$c = 6.41894$ (19) Å

$\beta = 90.795$ (3)°

$V = 1111.74$ (5) Å³

$Z = 8$

Mo $K\alpha$ radiation

$\mu = 0.42$ mm⁻¹

$T = 183$ K

$0.44 \times 0.17 \times 0.11$ mm

Data collection

Oxford Xcalibur Ruby CCD

diffractometer

Absorption correction: multi-scan

CrysAlis PRO (Oxford

Diffraction, 2009)

$T_{\min} = 0.891$, $T_{\max} = 0.955$

22537 measured reflections

5384 independent reflections

4400 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.033$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$

$wR(F^2) = 0.158$

$S = 1.21$

5384 reflections

181 parameters

16 restraints

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 0.82$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.69$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H1}\cdots\text{O7}^{\text{i}}$	0.78 (4)	1.80 (4)	2.570 (2)	168 (4)
$\text{O5}-\text{H5}\cdots\text{O8}^{\text{ii}}$	0.85 (4)	1.79 (4)	2.632 (2)	170 (4)
$\text{N11}-\text{H11A}\cdots\text{O4}^{\text{iii}}$	0.86 (2)	1.89 (2)	2.747 (2)	175 (4)
$\text{N11}-\text{H11B}\cdots\text{O8}^{\text{i}}$	0.86 (2)	2.10 (2)	2.951 (2)	171 (3)
$\text{N11}-\text{H11C}\cdots\text{O3}^{\text{iv}}$	0.88 (2)	1.99 (2)	2.852 (3)	169 (3)
$\text{N11}-\text{H11D}\cdots\text{O4}^{\text{i}}$	0.88 (2)	2.01 (2)	2.870 (2)	167 (3)
$\text{N12}-\text{H12A}\cdots\text{O6}^{\text{v}}$	0.87 (2)	1.91 (2)	2.755 (2)	165 (3)
$\text{N12}-\text{H12B}\cdots\text{O5}^{\text{vi}}$	0.88 (2)	2.16 (2)	3.008 (3)	161 (3)
$\text{N12}-\text{H12C}\cdots\text{O6}^{\text{i}}$	0.87 (2)	1.99 (2)	2.827 (2)	161 (3)
$\text{N12}-\text{H12D}\cdots\text{O2}$	0.88 (2)	1.88 (2)	2.754 (2)	175 (3)
$\text{N13}-\text{H13A}\cdots\text{O3}$	0.86 (2)	1.92 (2)	2.773 (2)	172 (3)
$\text{N13}-\text{H13B}\cdots\text{O2}^{\text{vii}}$	0.86 (2)	1.96 (2)	2.822 (2)	175 (3)
$\text{N13}-\text{H13C}\cdots\text{O6}^{\text{v}}$	0.89 (2)	1.95 (2)	2.830 (2)	168 (3)
$\text{N13}-\text{H13D}\cdots\text{O2}^{\text{ii}}$	0.86 (2)	1.96 (2)	2.820 (2)	176 (3)
$\text{N14}-\text{H14A}\cdots\text{O7}$	0.87 (2)	1.90 (2)	2.771 (2)	174 (3)
$\text{N14}-\text{H14B}\cdots\text{O8}^{\text{vii}}$	0.86 (2)	2.01 (2)	2.859 (2)	171 (3)
$\text{N14}-\text{H14C}\cdots\text{O3}^{\text{iv}}$	0.86 (2)	1.92 (2)	2.784 (2)	178 (3)
$\text{N14}-\text{H14D}\cdots\text{O4}$	0.89 (2)	1.89 (2)	2.771 (2)	171 (3)

Symmetry codes: (i) $-x, -y + 1, -z + 1$; (ii) $x, -y + \frac{1}{2}, z + \frac{1}{2}$; (iii) $-x, y + \frac{1}{2}, -z + \frac{3}{2}$; (iv) $-x, -y + 1, -z + 2$; (v) $x + 1, y, z + 1$; (vi) $x + 1, y, z$; (vii) $x, y, z + 1$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2009); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

BS thanks the University of Zürich for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BR2137).

References

- Akbayeva, D., Vaira, M., Costantini, S., Peruzzini, M. & Stoppioni, P. (2006). *Dalton Trans.* pp. 389–395.
- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Deifel, N. P., Holman, K. T. & Cahil, C. (2008). *Chem. Commun.* pp. 6037–6038.

- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Fernandez-Galan, R., Manzano, B., Otero, A., Lanfranchi, M. & Pellinghelli, M. (1994). *Inorg. Chem.* **33**, 2309–2312.
- Gebala, A. & Jones, M. (1969). *J. Inorg. Nucl. Chem.* **31**, 771–776.
- Khan, A. A., Roux, J. P. & James, W. J. (1972). *Acta Cryst.* **B28**, 2065–2069.
- Nikitenko, S., Berthon, C. & Moisy, P. (2007). *C. R. Chim.* **10**, 1122–1127.
- Oxford Diffraction (2009). *CrysAlis PRO*. Oxford Diffraction, Yarnton, England.
- Plakhotnyk, A., Ernst, L. & Schmutzler, R. (2005). *J. Fluorine Chem.* **126**, 27–31.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2010). E66, i26-i27 [doi:10.1107/S1600536810009839]

A new crystal modification of diammonium hydrogen phosphate, $(\text{NH}_4)_2(\text{HPO}_4)$

P. C. Kunz, C. Wetzel and B. Spingler

Comment

The addition of hexafluorophosphate salts (ammonium, silver, thallium or potassium) is usually used to precipitate complex cations from aqueous solutions. It is long known, that PF_6^- is sensitive towards hydrolysis under acidic conditions (Gebala and Jones 1969; Plakhotnyk, Ernst *et al.* 2005). In organic solvents the spatial hydrolysis to intermediate species as HF, POF_3 and PO_2F_2^- is observed (Fernandez-Galan, Manzano *et al.* 1994; Akbayeva, Vaira *et al.* 2006; Nikitenko, Berthon *et al.* 2007) and under hydrothermal conditions this reaction is used for the formation of phosphate materials (Deifel, Holman *et al.* 2008).

During the course of our investigation into coinage metal complexes of diphosphine ligands we used ammonium hexafluorophosphate in order to crystallise $[\text{Ag}(\text{diphosphine})_2]\text{PF}_6$ complexes. From these solutions we always obtained needle-like crystals which turned out to be the title compound $(\text{NH}_4)_2(\text{HPO}_4)$ as the product of hydrolysis of NH_4PF_6 .

A modification of diammonium hydrogen phosphate is known with the cell parameters $a = 11.043$ (6), $b = 6.700$ (3), $c = 8.031$ (4), $\beta = 113.42$ (3)° and $Z = 4$. (Khan, Roux *et al.* 1972) In the new modification of the title compound reported here, there are eight molecules of $(\text{NH}_4)_2(\text{HPO}_4)$ in the unit cell. The structure consists of PO_4 and NH_4 tetrahedra, held together by $\text{O}—\text{H}\cdots\text{O}$ and $\text{N}—\text{H}\cdots\text{O}$ hydrogen bonds. These hydrogen bonds are more or less linear ($169^\circ < \angle(\text{DHA}) < 178^\circ$). Of the four P—O bonds in both PO_4 tetrahedra, one is longer than the remaining three, typical of a $\text{O}_3\text{P}(\text{OH})$ group. The two different HPO_4^- molecules (P1O_4 and P2O_4) are hydrogen bonded to ten and seven ammonium ions, respectively. In hydrogen phosphate P2, one NH_4^+ molecule is bound to O5 and O7, two to O8 and three to O6. In the other hydrogen phosphate, the three non-protonated O-atoms O2, O3 and O4 are bound to three NH_4^+ molecules each, whereas the protonated O1 is only bound to one NH_4^+ molecule. In the hydrogen phosphate P1 the hydroxyl group O1H1 forms a hydrogen bridge to O7 ($d = 1.80$ (4) Å) and in the hydrogen phosphate P2 the hydroxyl group O5H5 forms a hydrogen bridge to O8 ($d = 1.79$ (4) Å). The N—O distances around the four-coordinated ammonium ions N13 and N14 are within the range of $2.77 < d < 2.86$ Å; The N \cdots O distances around N12 fall within this range with the exception of N12 \cdots O5 which is significantly longer ($d_{\text{N12O5}} = 3.007$ Å). A very different picture is found around N11. Here, five neighbouring O-atoms are found, three of which are in a shorter distance ($2.74 < d < 2.87$ Å) and two in a longer distance ($d_{\text{N11O5}} = 2.951$ Å and $d_{\text{N11O1}} = 3.048$ Å). The fifth N \cdots O contact may be a result of dynamic or static disorder of the ammonium ion or that each N atom in addition to three normal N—H \cdots O bonds also formed one bifurcated bond. This is in contrast to the other modification, in which each ammonium ion has five N \cdots O contacts smaller than 3.4 Å. Since Khan and Roux (Khan, Roux *et al.* 1972) only reported that they used "a commercially supplied crystalline sample", we cannot compare the crystallization conditions that lead to the two different crystal forms.

Experimental

A suitable crystal was covered with oil (Infinitec V8512, formerly known as Paratone N), mounted on top of a glass fibre and immediately transferred to the diffractometer. The final model was checked for higher symmetry with help of the program PLATON (Spek, 2009).

Refinement

All hydrogen atoms were located by difference Fourier synthesis and refined with fixed individual displacement parameters [$U(H) = 1.5U_{eq}(O)$] at an O-H distance of 0.87 Å.

Figures





Fig. 1. H-bonded network in the solid-state of the title compound. Displacement ellipsoids are drawn at a 50 % level, H-atoms are represented as capped sticks.

diammonium hydrogen phosphate

Crystal data



$$M_r = 132.06$$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$$a = 11.2868 (3) \text{ \AA}$$

$$b = 15.3466 (4) \text{ \AA}$$

$$c = 6.41894 (19) \text{ \AA}$$

$$\beta = 90.795 (3)^\circ$$

$$V = 1111.74 (5) \text{ \AA}^3$$

$$Z = 8$$

$$F(000) = 560$$

$$D_x = 1.578 \text{ Mg m}^{-3}$$

$$\text{Mo K}\alpha \text{ radiation, } \lambda = 0.7107 \text{ \AA}$$

Cell parameters from 11185 reflections

$$\theta = 2.7\text{--}37.6^\circ$$

$$\mu = 0.42 \text{ mm}^{-1}$$

$$T = 183 \text{ K}$$

Needle, colourless

$$0.44 \times 0.17 \times 0.11 \text{ mm}$$

Data collection

Oxford Xcalibur Ruby CCD
diffractometer

Radiation source: Enhance (Mo) X-ray Source
graphite

Detector resolution: 10.4498 pixels mm^{-1}

ω oscillation scan

Absorption correction: multi-scan
CrysAlis PRO (Oxford Diffraction, 2009)

$$T_{\min} = 0.891, T_{\max} = 0.955$$

5384 independent reflections

4400 reflections with $I > 2\sigma(I)$

$$R_{\text{int}} = 0.033$$

$$\theta_{\max} = 36.3^\circ, \theta_{\min} = 2.7^\circ$$

$$h = -18 \rightarrow 18$$

$$k = -25 \rightarrow 25$$

$$l = -9 \rightarrow 10$$

22537 measured reflections

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.048$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.158$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.21$	$w = 1/[\sigma^2(F_o^2) + (0.0508P)^2 + 1.9491P]$
5384 reflections	where $P = (F_o^2 + 2F_c^2)/3$
181 parameters	$(\Delta/\sigma)_{\max} < 0.001$
16 restraints	$\Delta\rho_{\max} = 0.82 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.69 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.12341 (16)	0.48227 (11)	0.6812 (3)	0.0260 (4)
H1	0.181 (4)	0.502 (3)	0.635 (6)	0.039*
O2	0.25104 (13)	0.34866 (10)	0.7264 (2)	0.0180 (3)
O3	0.13920 (15)	0.40992 (11)	1.0319 (2)	0.0198 (3)
O4	0.02645 (13)	0.34042 (10)	0.7334 (3)	0.0190 (3)
O5	-0.41866 (14)	0.29289 (11)	0.4731 (3)	0.0209 (3)
H5	-0.368 (3)	0.259 (3)	0.529 (6)	0.031*
O6	-0.47633 (13)	0.40326 (10)	0.2142 (2)	0.0174 (3)
O7	-0.30247 (15)	0.43252 (11)	0.4549 (2)	0.0209 (3)
O8	-0.28270 (13)	0.32102 (10)	0.1683 (2)	0.0182 (3)
P1	0.13662 (4)	0.39195 (3)	0.79806 (7)	0.01228 (10)
P2	-0.36645 (4)	0.36515 (3)	0.32105 (7)	0.01211 (10)
N11	0.03108 (16)	0.66864 (12)	0.7030 (3)	0.0185 (3)
H11A	0.009 (3)	0.7219 (13)	0.720 (5)	0.028*
H11B	0.1058 (17)	0.667 (2)	0.730 (5)	0.028*
H11C	-0.017 (3)	0.638 (2)	0.780 (5)	0.028*

supplementary materials

H11D	0.018 (3)	0.658 (2)	0.570 (3)	0.028*
N12	0.47645 (17)	0.41255 (13)	0.7925 (3)	0.0188 (3)
H12A	0.500 (3)	0.402 (2)	0.920 (3)	0.028*
H12B	0.523 (3)	0.382 (2)	0.711 (5)	0.028*
H12C	0.479 (3)	0.4675 (13)	0.760 (5)	0.028*
H12D	0.4031 (19)	0.395 (2)	0.770 (5)	0.028*
N13	0.29995 (16)	0.32905 (12)	1.2991 (3)	0.0182 (3)
H13A	0.252 (3)	0.351 (2)	1.207 (4)	0.027*
H13B	0.285 (3)	0.339 (2)	1.428 (3)	0.027*
H13C	0.3748 (18)	0.345 (2)	1.278 (5)	0.027*
H13D	0.288 (3)	0.2742 (12)	1.282 (5)	0.027*
N14	−0.18493 (17)	0.42049 (12)	0.8356 (3)	0.0178 (3)
H14A	−0.227 (3)	0.425 (2)	0.720 (4)	0.027*
H14B	−0.221 (3)	0.394 (2)	0.933 (4)	0.027*
H14C	−0.169 (3)	0.4726 (14)	0.877 (5)	0.027*
H14D	−0.1132 (19)	0.398 (2)	0.814 (5)	0.027*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0185 (7)	0.0193 (7)	0.0403 (10)	0.0021 (6)	0.0007 (6)	0.0145 (7)
O2	0.0140 (6)	0.0182 (6)	0.0220 (7)	0.0022 (5)	0.0025 (5)	−0.0015 (5)
O3	0.0225 (7)	0.0215 (7)	0.0153 (6)	0.0000 (5)	−0.0005 (5)	−0.0035 (5)
O4	0.0142 (6)	0.0179 (6)	0.0249 (7)	−0.0037 (5)	−0.0022 (5)	−0.0034 (5)
O5	0.0183 (6)	0.0207 (7)	0.0237 (7)	0.0025 (5)	0.0055 (5)	0.0099 (5)
O6	0.0145 (6)	0.0189 (6)	0.0188 (6)	0.0015 (5)	−0.0032 (5)	0.0032 (5)
O7	0.0221 (7)	0.0242 (7)	0.0165 (6)	−0.0048 (6)	−0.0035 (5)	−0.0031 (5)
O8	0.0150 (6)	0.0206 (7)	0.0192 (6)	−0.0010 (5)	0.0046 (5)	−0.0033 (5)
P1	0.01141 (19)	0.01162 (19)	0.01380 (19)	−0.00003 (14)	−0.00053 (14)	−0.00007 (14)
P2	0.01113 (18)	0.0140 (2)	0.01122 (18)	−0.00061 (14)	0.00038 (13)	0.00073 (14)
N11	0.0167 (7)	0.0184 (7)	0.0205 (8)	0.0015 (6)	−0.0007 (6)	−0.0001 (6)
N12	0.0170 (7)	0.0226 (8)	0.0169 (7)	0.0013 (6)	−0.0008 (6)	−0.0018 (6)
N13	0.0160 (7)	0.0199 (7)	0.0187 (7)	−0.0015 (6)	−0.0001 (5)	0.0002 (6)
N14	0.0178 (7)	0.0197 (7)	0.0159 (7)	−0.0006 (6)	0.0001 (5)	−0.0016 (6)

Geometric parameters (\AA , $^\circ$)

O1—P1	1.5821 (17)	N11—H11D	0.877 (18)
O1—H1	0.78 (4)	N12—H12A	0.869 (18)
O2—P1	1.5287 (15)	N12—H12B	0.878 (18)
O3—P1	1.5257 (16)	N12—H12C	0.868 (18)
O4—P1	1.5262 (15)	N12—H12D	0.879 (18)
O5—P2	1.5955 (16)	N13—H13A	0.863 (18)
O5—H5	0.85 (4)	N13—H13B	0.862 (18)
O6—P2	1.5254 (15)	N13—H13C	0.892 (18)
O7—P2	1.5201 (16)	N13—H13D	0.859 (18)
O8—P2	1.5295 (15)	N14—H14A	0.874 (18)
N11—H11A	0.862 (18)	N14—H14B	0.856 (18)
N11—H11B	0.859 (18)	N14—H14C	0.861 (18)

N11—H11C	0.876 (18)	N14—H14D	0.890 (18)
P1—O1—H1	116 (3)	H11C—N11—H11D	110 (3)
P2—O5—H5	115 (3)	H12A—N12—H12B	107 (3)
O3—P1—O4	111.44 (9)	H12A—N12—H12C	113 (3)
O3—P1—O2	111.70 (9)	H12B—N12—H12C	111 (3)
O4—P1—O2	112.43 (9)	H12A—N12—H12D	112 (3)
O3—P1—O1	107.95 (10)	H12B—N12—H12D	108 (3)
O4—P1—O1	104.71 (10)	H12C—N12—H12D	106 (3)
O2—P1—O1	108.21 (9)	H13A—N13—H13B	118 (3)
O7—P2—O6	111.73 (9)	H13A—N13—H13C	112 (3)
O7—P2—O8	111.74 (9)	H13B—N13—H13C	107 (3)
O6—P2—O8	112.79 (9)	H13A—N13—H13D	101 (3)
O7—P2—O5	107.68 (9)	H13B—N13—H13D	105 (3)
O6—P2—O5	103.69 (9)	H13C—N13—H13D	113 (3)
O8—P2—O5	108.73 (9)	H14A—N14—H14B	114 (3)
H11A—N11—H11B	107 (3)	H14A—N14—H14C	107 (3)
H11A—N11—H11C	105 (3)	H14B—N14—H14C	109 (3)
H11B—N11—H11C	119 (3)	H14A—N14—H14D	112 (3)
H11A—N11—H11D	105 (3)	H14B—N14—H14D	112 (3)
H11B—N11—H11D	110 (3)	H14C—N14—H14D	102 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1 \cdots O7 ⁱ	0.78 (4)	1.80 (4)	2.570 (2)	168 (4)
O5—H5 \cdots O8 ⁱⁱ	0.85 (4)	1.79 (4)	2.632 (2)	170 (4)
N11—H11A \cdots O4 ⁱⁱⁱ	0.86 (2)	1.89 (2)	2.747 (2)	175 (4)
N11—H11B \cdots O8 ⁱ	0.86 (2)	2.10 (2)	2.951 (2)	171 (3)
N11—H11C \cdots O3 ^{iv}	0.88 (2)	1.99 (2)	2.852 (3)	169 (3)
N11—H11D \cdots O4 ⁱ	0.88 (2)	2.01 (2)	2.870 (2)	167 (3)
N12—H12A \cdots O6 ^v	0.87 (2)	1.91 (2)	2.755 (2)	165 (3)
N12—H12B \cdots O5 ^{vi}	0.88 (2)	2.16 (2)	3.008 (3)	161 (3)
N12—H12C \cdots O6 ⁱ	0.87 (2)	1.99 (2)	2.827 (2)	161 (3)
N12—H12D \cdots O2	0.88 (2)	1.88 (2)	2.754 (2)	175 (3)
N13—H13A \cdots O3	0.86 (2)	1.92 (2)	2.773 (2)	172 (3)
N13—H13B \cdots O2 ^{vii}	0.86 (2)	1.96 (2)	2.822 (2)	175 (3)
N13—H13C \cdots O6 ^v	0.89 (2)	1.95 (2)	2.830 (2)	168 (3)
N13—H13D \cdots O2 ⁱⁱ	0.86 (2)	1.96 (2)	2.820 (2)	176 (3)
N14—H14A \cdots O7	0.87 (2)	1.90 (2)	2.771 (2)	174 (3)
N14—H14B \cdots O8 ^{vii}	0.86 (2)	2.01 (2)	2.859 (2)	171 (3)
N14—H14C \cdots O3 ^{iv}	0.86 (2)	1.92 (2)	2.784 (2)	178 (3)
N14—H14D \cdots O4	0.89 (2)	1.89 (2)	2.771 (2)	171 (3)

Symmetry codes: (i) $-x, -y+1, -z+1$; (ii) $x, -y+1/2, z+1/2$; (iii) $-x, y+1/2, -z+3/2$; (iv) $-x, -y+1, -z+2$; (v) $x+1, y, z+1$; (vi) $x+1, y, z$; (vii) $x, y, z+1$.

